

BROMHEXIN® BC *Menarini*

GENERIC NAME

Bromhexine

COMPOSITION

Bromhexin® 4 Berlin-Chemie: 5 ml solution (1 measuring spoon/measuring cap) contains Bromhexine hydrochloride 4 mg, propylene glycol 1.25 g, sorbitol 2.0 g (0.17 bread exchange).

Bromhexin® 8 Berlin-Chemie: 5 ml solution (1 measuring spoon/measuring cap) contains Bromhexine hydrochloride 8 mg, propylene glycol 1.25 g, sorbitol 2.0 g (0.17 bread exchange).

Bromhexin® 8 Berlin-Chemie: 1 coated tablet contains bromhexine hydrochloride 8 mg.

Bromhexin® 10 Berlin-Chemie: 5 ml solution (1 measuring spoon/measuring cap) contains Bromhexine hydrochloride 10 mg, propylene glycol 1.25 g, sorbitol 2.0 g (0.17 bread exchange).

Bromhexin® 12 Berlin-Chemie: 1 ml solution (25 drops) contains Bromhexine hydrochloride 12 mg.

INDICATIONS

Secretolytic treatment of acute and chronic broncho pulmonary diseases accompanied by impaired formation and transport of mucus.

DOSAGE AND ADMINISTRATION

Children (under 6 years): 4 mg bromhexine hydrochloride t.i.d.

Children (6-14 years) and patients weighting less than 50 kg: 8 mg bromhexine hydrochloride t.i.d.

Adolescent (over 14 years) and adults: 8-16 mg bromhexine hydrochloride t.i.d.

SHELF-LIFE/STORAGE INSTRUCTIONS

Shelf-life: Bromhexin Berlin-Chemie 8 (coated tablets): 5 years.

Bromhexin Berlin-Chemie 4/8/10/12 (solution): 3 years.

PACKAGE

Bromhexin® 4 Berlin-Chemie: bottle of 100 ml solution.

Bromhexin® 8 Berlin-Chemie: bottle of 100 ml solution, box of 50 coated tablets.

Bromhexin® 10 Berlin-Chemie: bottle of 100 ml solution.

Bromhexin® 12 Berlin-Chemie: bottle of 50 ml solution (drops).